



# Myocarditis and Pericarditis After mRNA COVID-19 Vaccination

Updated June 19, 2022

CDC and its partners are actively monitoring reports of myocarditis and pericarditis after COVID-19 vaccination. Active monitoring includes reviewing data and medical records and evaluating the relationship to COVID-19 vaccination.

**Myocarditis** is inflammation of the heart muscle, and **pericarditis** is inflammation of the outer lining of the heart. In both cases, the body's immune system causes inflammation in response to an infection or some other trigger. Learn more about [myocarditis and pericarditis](#). [🔗](#) **Seek medical care if you or your child have symptoms of these conditions after COVID-19 vaccination.** Myocarditis and pericarditis have rarely been reported, especially in adolescents and young adult males within several days after COVID-19 vaccination.

## What You Need to Know

- Cases of myocarditis reported to the [Vaccine Adverse Event Reporting System \(VAERS\)](#) [🔗](#) have occurred:
  - After mRNA COVID-19 vaccination (Pfizer-BioNTech or Moderna), especially in male adolescents and young adults
  - More often after the second dose
  - Usually within a week of vaccination
- Most patients with myocarditis or pericarditis who received care responded well to medicine and rest and felt better quickly.
- Patients can usually return to their normal daily activities after their symptoms improve. Those who have been diagnosed with myocarditis should consult with their cardiologist (heart doctor) about return to exercise or sports. More information will be shared as it becomes available.

Both myocarditis and pericarditis have the following symptoms:

- Chest pain
- Shortness of breath
- Feelings of having a fast-beating, fluttering, or pounding heart

Seek medical care if you or your child have any of the specific or general symptoms of myocarditis or pericarditis especially if it's within a week after COVID-19 vaccination.

If you have any health problems after vaccination, report them to [VAERS](#) [🔗](#).

Healthcare Providers: For additional recommendations and clinical guidance, visit [Clinical Considerations: Myocarditis after mRNA COVID-19 Vaccines](#).

## Should I Still Get Myself or My Child Vaccinated?

Yes. **CDC continues to recommend that everyone ages 6 months and older get vaccinated for COVID-19.** The known risks of COVID-19 illness and its related, possibly severe complications, such as long-term health problems, hospitalization, and even death, far outweigh the potential risks of having a rare adverse reaction to vaccination, including the possible risk of myocarditis or pericarditis.

If you have already gotten the first dose of Pfizer-BioNTech or Moderna vaccine, or if your child has already gotten the first dose of the Pfizer-BioNTech vaccine, it's important to get the [second dose](#) unless a vaccination provider or your doctor tells you [not to get it](#).

If you have concerns about COVID-19 vaccination, talk with your healthcare provider or your child's doctor, nurse, or clinic.

Last Updated June 19, 2022